

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/50		A2	(11) International Publication Number: WO 99/39689
			(43) International Publication Date: 12 August 1999 (12.08.99)
(21) International Application Number: PCT/EP99/00320 (22) International Filing Date: 19 January 1999 (19.01.99) (30) Priority Data: 09/017,746 3 February 1998 (03.02.98) US (71) Applicant (for AU BB CA CY GB GD GH GM IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZW only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). (71) Applicant (for all designated States except AU BB CA CY GB GD GH GM IE IL IN KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZW): UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL). (71) Applicant (for IN only): HINDUSTAN LEVER LIMITED [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Mumbai 400 020, Maharashtra (IN).		(72) Inventors: VINSKI, Paul; Chesebrough Pond's USA Co., 40 Merritt Boulevard, Trumbull, CT 06611 (US). ZNAIDEN, Alexander, Paul; Chesebrough Pond's USA Co., 40 Merritt Boulevard, Trumbull, CT 06611 (US). SLAVTCHEFF, Craig, Stephen; Chesebrough Pond's USA Co., 40 Merritt Boulevard, Trumbull, CT 06611 (US). (74) Agent: ROTS, Maria, Johanna, Francisca; Unilever plc, Patent Dept., Colworth House, Sharnbrook, Bedford MK44 1LQ (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.	
(54) Title: FOAMING CLEANSING SKIN PRODUCT			
<div style="text-align: center;"> $R-C(=O)-NH-C_2H_4-N(C_2H_4OR^1)-C_2H_4R^2$ </div> <div style="text-align: right;">(I)</div>			
(57) Abstract			
<p>A foaming cleansing product is provided as a cleansing composition packaged in a non-aerosol mechanical dispenser. The dispenser includes a container for holding a liquid composition, a dispensing head with a housing enclosing a pump mechanism and a screen material in the flow path to convert liquid composition into a foam, and a diptube for delivering liquid from the container to the dispensing head. The cleansing composition is substantially free of water insoluble emollients and includes an anionic surfactant and an amphoteric surfactant of structure (I); wherein R is a fatty alkyl of 6 to 22 carbons; R¹ is selected from the group consisting of H, C₂H₄COOH and C₂H₄COONa; and R² is selected from the group consisting of COONa, CH₂CHOHCH₂SO₃Na, C₂H₄COONa, CH₂COOCH₂COOH and CH₂COOCH₂COONa.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

FOAMING CLEANSING SKIN PRODUCT

BACKGROUND OF THE INVENTION

5 Field of the Invention

The invention concerns a foaming cleansing product which is a combination of a non-aerosol dispenser and a cleansing composition designed to cooperate with the dispenser for
10 generating a mousse quality foam while concurrently imparting moisturization benefits to the skin.

The Related Art

15 Cleansing compositions in mousse form have certain appeal to consumers. Foremost is the instant foam achieved by the mere press of a button. Aerosol dispensers employing propellants generally provide a satisfactory foam volume. Unfortunately, aerosol products are under attack for
20 environmental reasons. Volatile organics interfere with the earth's ozone layer and contribute to smog in metropolitan areas. Aerosol packages are also relatively costly to assemble. For all these reasons, attention has been recently directed at non-aerosol dispensers.

25

U.S. Patent 5,635,469 (Fowler et al.) discloses personal cleansing products comprising a foamable liquid composition and a foam-producing non-aerosol dispenser. The compositions include a surfactant, a water soluble cationic
30 or nonionic polymer, a humectant, a water-insoluble

- 2 -

emollient and water. The dispenser employs at least two screens through which the composition is blown to generate a foam.

5 Delivery of cleansing compositions via non-aerosol dispensers has presented many challenges. Foams produced from these dispensers often lack the dense volume consumers desire. These foams also may not be long lasting or
10 luxurious. Additives within these compositions intended to deliver skin benefit agents can interfere with the foam properties. Still further, certain types of non-aerosol dispensers which operate with porous filters or meshed screens require the cleansing product to be relatively non-viscous. Where the package is transparent, formulators
15 seek clear formulations for aesthetics purposes.

Accordingly, it is an object of the present invention to provide a cleansing product in mousse form which delivers a dense luxurious foam.

20

Another object of the present invention is to provide a cleansing product in mousse form which delivers benefit agents to the skin that do not interfere with foam properties.

25

Still another object of the present is to provide a cleansing product in mousse form having a transparent liquid formulation.

- 3 -


These and other objects of the present invention will become more readily apparent from consideration of the summary and detailed description which follows.

5 SUMMARY OF THE INVENTION

A foaming cleansing product is provided which includes:

- (A) a non-aerosol dispenser having:
- 10 (i) a container for storing a cleansing composition;
 - (ii) a head having a housing surrounding a pump mechanism and a foam-forming screen material;
 - (iii) a diptube communicating between the container and head functioning to fluidly deliver liquid
 - 15 cleansing composition between container and head and being upstream from the screen material; and
- (B) the cleansing composition being substantially free of
- 20 water insoluble emollients and including:
 - (i) from 0.1 to 50% by weight of an anionic surfactant; and
 - (ii) from 0.1 to 30% by weight of an amphoteric surfactant having the structure:

25




- 4 -

wherein R is a fatty alkyl of 6 to 22 carbons;
 R^1 is selected from the group consisting of H,
 C_2H_4COOH and C_2H_4COONa ; and R^2 is selected from
the group consisting of $COONa$, $CH_2CHOHCH_2SO_3Na$,
5 C_2H_4COONa , CH_2COOCH_2COOH and CH_2COOCH_2COONa .

DETAILED DESCRIPTION

Now it has been discovered that a rich, luxurious creamy
10 foam can be generated through a non-aerosol mechanical pump
by a cleansing system based on anionic surfactants and an
amphoteric surfactant having the structure:



wherein R is a fatty alkyl of 6 to 22 carbons; R^1 is
selected from the group consisting of H, C_2H_4COOH and
 C_2H_4COONa ; and R^2 is selected from the group consisting of
20 $COONa$, $CH_2CHOHCH_2SO_3Na$, C_2H_4COONa , CH_2COOCH_2COOH and
 CH_2COOCH_2COONa .

Specific examples of these amphoteric surfactants include
the alkali, alkaline earth, ammonium and trialkanolammonium
25 salts of cocoamphoacetate, cocoamphodiacetate,
cocoamphopropionate, cocoamphodipropionate and mixtures
thereof. Most preferred is sodium cocoamphoacetate
available as Miranol HMA from the Rhone Poulenc Corporation.

- 5 -

Similar surfactants are also available as Amphoterger® from Lonza Inc., Fair Lawn, New Jersey. While the sodium salt is preferred, other cations can also be employed including lithium, potassium, magnesium and calcium. Amounts of the amphoteric surfactant may range from 0.1 to 20%, preferably from 1 to 10%, optimally from 2 to 6% by weight of the composition.

A further component of cleansing compositions according to the present invention is that of an anionic surfactant. Illustrative but not limiting examples include the following classes:

- (1) Alkyl benzene sulfonates in which the alkyl group contains from 9 to 15 carbon atoms, preferably 11 to 14 carbon atoms in straight chain or branched chain configuration. Especially preferred is a linear alkyl benzene sulfonate containing about 12 carbon atoms in the alkyl chain.
- (2) Alkyl sulfates obtained by sulfating an alcohol having 8 to 22 carbon atoms, preferably 12 to 16 carbon atoms. The alkyl sulfates have the formula $\text{ROSO}_3^- \text{M}^+$ where R is the C₈₋₂₂ alkyl group and M is a mono- and/or divalent cation.
- (3) Paraffin sulfonates having 8 to 22 carbon atoms, preferably 12 to 16 carbon atoms, in the alkyl moiety. These surfactants are commercially available as Hostapur SAS from Hoechst Celanese.

- 6 -

- (4) Olefin sulfonates having 8 to 22 carbon atoms, preferably 12 to 16 carbon atoms. Most preferred is sodium C₁₄-C₁₆ olefin sulfonate, available as Bioterge AS 40®
- 5
- (5) Alkyl ether sulfates derived from an alcohol having 8 to 22 carbon atoms, preferably 12 to 16 carbon atoms, ethoxylated with less than 30, preferably less than 12, moles of ethylene oxide. Most preferred is sodium lauryl ether sulfate formed from 2 moles average ethoxylation, commercially available as Standopol ES-2®.
- 10
- (6) Alkyl glyceryl ether sulfonates having 8 to 22 carbon atoms, preferably 12 to 16 carbon atoms, in the alkyl moiety.
- 15
- (7) Fatty acid ester sulfonates of the formula: $R^1CH(SO_3-M)CO_2R^2$ where R^1 is straight or branched alkyl from about C₈ to C₁₈, preferably C₁₂ to C₁₆, and R^2 is straight or branched alkyl from about C₁ to C₆, preferably primarily C₁, and M+ represents a mono- or divalent cation.
- 20
- (8) Secondary alcohol sulfates having 6 to 18, preferably 8 to 16 carbon atoms.
- 25
- (9) Fatty acyl isethionates having from 10 to 22 carbon atoms, with sodium cocoyl isethionate being preferred.

- 7 -

(10) Dialkyl sulfosuccinates wherein the alkyl groups range from 3 to 20 carbon atoms each.

5 (11) Alkanoyl sarcosinates corresponding to the formula
RCON(CH₃)CH₂CH₂CO₂M wherein R is alkyl or alkenyl of
about 10 to about 20 carbon atoms and M is a water-
soluble cation such as ammonium, sodium, potassium and
trialkanolammonium. Most preferred is sodium lauroyl
10 sarcosinate.

Co-surfactants may also be present to aid in the foaming,
detergency and mildness properties. Nonionic and amphoteric
actives are the preferred co-surfactants. Suitable nonionic
15 surfactants include C₁₀-C₂₀ fatty alcohol or acid hydrophobes
condensed with from 2 to 100 moles of ethylene oxide or
propylene oxide per mole of hydrophobe; C₂-C₁₀ alkyl phenols
condensed with from 2 to 20 moles of alkylene oxides; mono-
and di- fatty acid esters of ethylene glycol such as
20 ethylene glycol distearate; fatty acid monoglycerides;
sorbitan mono- and di- C₈-C₂₀ fatty acids; and
polyoxyethylene sorbitan available as Polysorbate 80 and
Tween 80® as well as combinations of any of the above
surfactants.

25

Other useful nonionic surfactants include alkyl
polyglycosides, saccharide fatty amides (e.g. methyl
gluconamides) as well as long chain tertiary amine oxides.
Examples of the latter category are: dimethyldodecylamine

- 8 -

oxide, oleyldi(2-hydroxyethyl)amine oxide,
dimethyloctylamine oxide, dimethyldecylamine oxide,
dimethyltetradecylamine oxide, di(2-
hydroxyethyl)tetradecylamine oxide, 3-didodecoxy-2-
5 hydroxypropyldi(3-hydroxypropyl)amine oxide, and
dimethylhexadecylamine oxide.

Amounts of the nonionic surfactant may range from 0.1 to
40%, preferably from 0.5 to 15%, optimally from 1 to 5% by
10 weight of the total composition.

Amphoteric surfactants such as betaines may also be employed
as co-actives along with the anionic surfactants. Suitable
betaines may have the general formula $RN^+(R^1)_2R^2COO^-$ wherein
15 R is a hydrophobic moiety selected from the group consisting
of alkyl groups containing from 10 to 22 carbon atoms,
preferably from 12 to 18 carbon atoms; alkyl aryl and aryl
alkyl groups containing 10 to 22 carbon atoms with a benzene
ring being treated as equivalent to about 2 carbon atoms,
20 and similar structures interrupted by amido or ether
linkages; each R^1 is an alkyl group containing from 1 to 3
carbon atoms; and R^2 is an alkylene group containing from 1
to about 6 carbon atoms. Sulfobetaines such as
cocoamidopropyl sultaine are also suitable.

25 Examples of preferred betaines are dodecyl dimethyl betaine,
cetyl dimethyl betaine, dodecyl amidopropyldimethyl betaine,
tetradecyldimethyl betaine, tetradecylamidopropyldimethyl
betaine, and dodecyldimethylammonium hexanoate. Most
30 preferred is cocoamidopropyl betaine available as

- 9 -

Tegobetaine F® sold by Th. Goldschmidt AG of Germany. Amounts of the betaine may range from 0.05 to 15%, preferably from 0.5 to 10%, optimally from 2 to 8% by weight of the total composition.

5

Moisturizing ingredients may also be included in compositions of the present invention. Water soluble moisturizers such as polyhydric alcohol-type humectants are particularly preferred. Typical polyhydric alcohols include
10 glycerol (also known as glycerin), polyalkylene glycols and more preferably alkylene polyols and their derivatives, including propylene glycol, dipropylene glycol, polypropylene glycol, polyethylene glycol and derivatives thereof, sorbitol, hydroxypropyl sorbitol, hexylene glycol,
15 1,3-butylene glycol, 1,2,6-hexanetriol, ethoxylated glycerol, propoxylated glycerol and mixtures thereof. For best results the humectant is preferably glycerin. The amount of humectant may range anywhere from 0.5 to 30%, preferably between 1 and 15% by weight of the composition.

20

Compositions of the present invention preferably are substantially free of any oil phase, especially substantially free of water insoluble emollients. The term "substantially free" means less than 0.05%, preferably less
25 than 0.01% emollient, and water insoluble means any emollient having a solubility in distilled water at 25°C of less than about 1 gm per 100 mL, more preferably less than about 0.1 gm per 100 mL. Absent water insoluble emollients, the compositions can be transparent and have improved
30 foamability.

- 10 -

Preservatives may be incorporated into the cosmetic compositions of this invention to protect against the growth of potentially harmful microorganisms. Suitable traditional preservatives are EDTA salts and alkyl esters of para-
5 hydroxybenzoic acid. Other preservatives which have more recently come into use include hydantoin derivatives, propionate salts, and a variety of quaternary ammonium compounds. Cosmetic chemists are familiar with appropriate preservatives and routinely choose them to satisfy the
10 preservative challenge test and to provide product stability. Particularly preferred preservatives are iodopropynyl butyl carbamate, phenoxyethanol, methyl paraben, propyl paraben, imidazolidinyl urea, sodium dehydroacetate and benzyl alcohol. The preservatives should
15 be selected having regard for the use of the composition and possible incompatibilities between the preservatives and other ingredients in the composition. Preservatives are preferably employed in amounts ranging from 0.01% to 2% by weight of the composition.

20 Minor adjunct ingredients may be present in the cosmetic compositions. Among them may be the water-soluble vitamins, colorants, fragrances and opacifiers. Each of these substances may range from 0.05 to 5%, preferably between 0.1
25 and 3% by weight of the composition.

Advantageously, the compositions of the invention may contain a foam densifying agent. Examples of this substance are waxy materials with a melting point greater than 20°C,
30 preferably greater than 40°C. Illustrative are ethoxylated

- 11 -


glyceride esters such as PEG 75 soy glycerides sold under the trademark Acconon S 75. Also useful are C₈-C₁₂ acyl lactylates such as sodium lauroyl lactylate sold as Pationic 138 C[®] available from the Patterson Chemical Company.

- 5 Amounts of these agents may range from 0.1 to 10%, preferably from 0.5 to 5%, optimally from 1 to 3% by weight.

- Cationic conditioning agents in monomeric and polymeric type are also useful for purposes of this invention. Examples of the polymeric type include: cationic cellulose derivatives, cationic starches, copolymers of a diallyl quaternary ammonium salt and an acryl amide, quaternized vinylpyrrolidone vinylimidazole polymers polyglycol amine condensates, quaternized collagen polypeptide, polyethylene imine, cationized silicon polymer (e.g. Amodimethicone), cationic silicon polymers provided in a mixture with other components under the trademark Dow Corning 929 (cationized emulsion), copolymers of adipic acid and dimethylaminohydroxypropyl diethylenetriamine, cationic chitin derivatives, cationized guar gum (e.g. Jaguar C-B-S, Jaguar C-17, Jaguar C-16, etc. manufactured by the Celanese Company), quaternary ammonium salt polymers (e.g. Mirapol A-15, Mirapol AD-1, Mirapol AZ-1, etc., manufactured by the Miranol Divison of the Rhone Poulenc Company). Most preferred is polyquaternium-11 available as Luviquat[®] PQ 11 sold by the BASF Corporation.

Examples of monomeric cationic conditioning agents are salts of the general structure:

- 12 -



5

wherein R^1 is selected from an alkyl group having from 12 to 22 carbon atoms, or aromatic, aryl or alkaryl groups having from 12 to 22 carbon atoms; R^2 , R^3 , and R^4 are independently selected from hydrogen, an alkyl group having from 1 to 22 carbon atoms, or aromatic, aryl or alkaryl groups having from 12 to 22 carbon atoms; and X^- is an anion selected from chloride, bromide, iodide, acetate, phosphate, nitrate, sulfate, methyl sulfate, ethyl sulfate, tosylate, lactate, citrate, glycolate, and mixtures thereof. Additionally, the alkyl groups can also contain ether linkages, or hydroxy or amino group substituents (e.g., the alkyl groups can contain polyethylene glycol and polypropylene glycol moieties). Preferably the anion is phosphate, especially preferred is hydroxy ethyl cetyl dimonium phosphate available as Luviquat® Mono CP from the BASF Corporation.

Amino silicones quats may similarly be employed. Most preferred is Silquat AD designated by the CTFA as Silicone Quaternium 8, available from Siltech Inc.

Amounts of each cationic agent may range from 0.05 to 5%, preferably from 0.1 to 3%, optimally from 0.3 to 2.5% by weight.

30

- 13 -

Advantageously the compositions of this invention are transparent. By the term transparent is meant having a maximum transmittance of light of at least 4% of any wave length in the range of 400 to 700 nm through a sample 1 cm
5 thick. A transparent composition is one which also permits sufficient light transmittance to enable reading of newspaper print through a thickness commensurate with a diameter of the container employed with the herein described dispenser.

10

Compositions of this invention should also be of relatively low viscosity to be pumpable. Viscosity may range from 1 to 300 centipoise, preferably from 3 to 100 centipoise, optimally from 5 to 50 centipoise at 25°C.

15

An essential element of cleansing products according to this invention is a non-aerosol mechanical dispenser. The dispenser is generally characterized by a container for storing the composition (preferably a transparent
20 container), a dispensing head defined by a housing containing a pump, and a diptube for transferring the composition from the container into the dispensing head. Foam is created by requiring the composition to pass through a screen material which may be a porous substance such as a
25 sintered material, a wire (plastic or metal) gauze screen or similar structures.

Suitable dispensers are described in U.S. Patent 3,709,437 (Wright), U.S. Patent 3,937,364 (Wright), U.S. Patent
30 4,022,351 (Wright), U.S. Patent 4,147,306 (Bennett), U.S.

- 14 -

Patent 4,184,615 (Wright), U.S. Patent 4,598,862 (Rice),
U.S. Patent 4,615,467 (Grogan et al.) and U.S. Patent
5,364,031 (Tamiguchi et al.). Most preferred however is a
device owned by the Airspray International Corporation
5 described in WO 97/13585 (Van der Heijden). All these
patents are incorporated herein by reference. The Airspray
device comprises a container for storing a cleansing
composition and a dispensing head, the latter including at
least a concentric air pump and liquid pump. Each of the
10 pumps has a piston chamber with a piston displaceable
therein and an inlet and discharge, and an operating
component for operating the two pumps. The operating
component is integral with one of the pistons and comprises
an outflow channel with a dispensing opening. Shut-off
15 mechanisms, rendering it possible to suck up air or liquid,
respectively, and dispense them, are present in the inlet
and discharge of the pumps. The air pump includes a double-
acting shut-off device which can be operated actively by the
operating component. The shut-off device prevents both the
20 inlet of air to the air pump and discharge of air therefrom.
The air piston is able to be moved freely at least over a
small distance with respect to the operating component.

The following examples will more fully illustrate the
25 embodiments of this invention. All parts, percentages and
proportions referred to herein and in the
appended claims are by weight of the total composition
unless otherwise indicated.

- 15 -

EXAMPLES 1-8

The following examples are representative cleansing formulations according to the present invention.

5

[illegible]

- 17 -

These formulas are prepared by charging water to form Phase A in a main vessel. Heating is then begun to achieve a 50°C temperature. Pationic 138C® is then added and allowed to slowly dissolve. The remaining ingredients listed under
5 Phase A are added and allowed to dissolve. Heating and stirring are continued until a uniform mixture is obtained. Thereafter Phase A is cooled. Phase B ingredients are then added to Phase A with slow agitation to minimize aeration. Stirring is continued until the mixture becomes uniform.
10 Premix Phase C is then added to the resultant mixture at 35°C.

EXAMPLES 9-18

15 The following examples demonstrate the effect of changing surfactants and additives on the foam and skinfeel properties.

COMPONENT	EXAMPLE																	
	9	10	11	12	13	14	15	16	17	18								
Phase A																		
Paticonic 138C (sodium lauroyl lactylate)	--	--	--	--	--	--	1.00	--	1.00	--							--	
Acconon S75 (PEG 75 soy glycerides)	--	--	--	--	--	--	--	1.00	1.00	--							--	
Silquat AD	--	--	--	--	--	--	--	--	--	--							1.00	
Water	90.00	90.00	90.00	90.00	90.00	80.00	79.00	79.00	79.00	78.00							79.00	
Phase B																		
Standapol ES-2 (SLES 28%)	10.00	--	--	--	5.00	5.00	5.00	5.00	5.00	5.00							5.00	
Biorterge AS 40 (sodium C14-16 olefin sulphionate)	--	10.00	--	--	5.00	5.00	5.00	5.00	5.00	5.00							5.00	
Tegobetaine F (Cocoamidopropyl betaine)	--	--	10.00	--	--	--	--	--	--	--							--	
Miranol HMA (sodium lauroamphoacetate 30%)	--	--	--	10.00	--	10.00	10.00	10.00	10.00	10.00							10.00	

- 19 -

PERFORMANCE	
9	Good dispensing and foam build-up; lather very thin, lacks substance; very stripping afterfeel
10	Good dispensing and foam build-up; lather very thin, lacks substance; very stripping afterfeel
11	Good dispensing and foam build-up; lather very thin, poor cleansing; mild afterfeel
12	Good dispensing but foam collapses quickly; lather thin, poor cleansing; mild afterfeel; cloudy
13	Good dispensing and foam build-up; lather very thin, lacks substance, very stripping afterfeel
14	Good dispensing and foam build-up; lather good but could be denser; afterfeel not stripping
15	Good dispensing and foam build-up; lather good but could be denser; afterfeel better, more substantive; not stripping
16	Good dispensing and foam build-up; lather dense; afterfeel better, not stripping
17	Good dispensing and foam build-up; lather denser, afterfeel good, mild not stripping
18	Good dispensing and foam build-up; lather good but could be denser; afterfeel good, conditioned

The comparative experiments indicate that with but one exception all of these examples have good dispensing and foam build-up. However, use of anionic surfactants alone or even amphoteric surfactants alone as in Examples 9 through 13 results in a relatively very thin lather which lacks substance. These compositions also imparted a relatively harsh afterfeel with stripping of oils from the skin.

Example 14 employing a combination of anionic and amphoteric surfactant however provided a relatively non-stripping afterfeel; lather was also much denser than in the earlier examples. The addition of sodium lauroyl lactylate or PEG 75 soy glycerides as in Examples 15 through 18 substantially enhanced lather and provided very good afterfeel.

- 20 -

EXAMPLES 19-26

The following examples were prepared to investigate the effect of water-insoluble emollients on foam, afterfeel and other physical properties.

COMPONENT	EXAMPLE							
	19	20	21	22	23	24	25	26
Phase A								
Pationic 138C (sodium lauroyl lactylate)	--	1.00	--	1.00	1.00	1.00	1.00	1.00
Acconon S75 (PEG 75 soy glycerides)	--	--	1.00	1.00	1.00	1.00	1.00	1.00
Water	79.00	78.00	78.00	77.00	77.00	77.00	77.00	77.00
Silquat AD or emollient**	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Phase B								
Standapol ES-2 (SLES 28%)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00
Bioterge AS 40 (sodium C14-16 olefin sulphonate)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00
Miranol HMA (sodium lauroamphoacetate 30%)	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00

	PERFORMANCE
19	Good dispensing and foam build-up; lather good but could be denser; afterfeel good, conditioned
20	Good dispensing and foam build-up; lather good but could be denser; afterfeel good, more substantive, conditioned not stripped
21	Good dispensing and foam build-up; lather denser; afterfeel better, more conditioning
22	Good dispensing and foam build-up; lather denser; afterfeel excellent, conditioned and moisturized
23	Dimethicone instead of Silquat AD - dimethicone incompatible, separates into two phases
24	Dimethicone microemulsion instead of Silquat AD - disperses but depresses foam and lather build-up
25	Sunflower seed oil instead of Silquat AD - dispersed but depressed foam and lather; haze created
26	Maleated soybean oil instead of Silquat AD - separated into two phases

- 21 -

Examples 23-26 demonstrate that water insoluble emollients such as dimethicone, dimethicone microemulsions, sunflower seed oil and maleated soybean oil all either depress the lather or separated out from the formulation.

- 22 -

CLAIMS

1. A foaming cleansing product comprising:


5 (A) a non-aerosol dispenser having:

- (i) a container for storing a cleansing composition;
- 10 (ii) a head having a housing surrounding a pump mechanism and a foam-forming screen material;
- (iii) a diptube communicating between the container and head functioning to fluidly deliver liquid cleansing composition
- 15 between container and head and being upstream from the screen material; and

(B) the cleansing composition being substantially free of water insoluble emollients and comprising:

20

- (i) from 0.1 to 50% by weight of an anionic surfactant; and
- (ii) from 0.1 to 30% by weight of an amphoteric surfactant having structure:
- 25



- 23 -

- wherein R is a fatty alkyl of 6 to 22 carbons; R¹ is selected from the group consisting of H, C₂H₄COOH and C₂H₄COONa; and R₂ is selected from the group consisting of COONa, CH₂CHOHCH₂SO₃Na, C₂H₄COONa, CH₂COOCH₂COOH and CH₂COOCH₂COONa.
- 5
2. The product according to claim 1 wherein the screen material is a wire gauze.
- 10 3. The product according to claim 1 or 2 wherein a second wire gauze is placed downstream from the first wire gauze, the cleansing composition being required to traverse both the first and second wire gauze to achieve a foam.
- 15
4. The product according to any preceding claim wherein the amphoteric surfactant is selected from the group consisting of lauroamphoacetate and lauroamphodiacetate.
- 20
5. The product according to any preceding claim further comprising a foam densifying agent present in an amount from 0.1 to 10% by weight.
- 25 6. The product according to claim 5 wherein the densifying agent is an ester wax with a melting point greater than 20°C.

- 24 -

7. The product according to claim 6 wherein the ester wax has a melting point greater than 40°C.
8. The product according to claim 6 or 7 wherein the ester wax is an ethoxylated glyceride ester.
9. The product according to any preceding claim wherein the foam densifying agent is a C₈-C₁₂ acyl lactylate.
- 10 10. The product according to claim 9 wherein the lactylate is sodium lauroyl lactylate.
11. The product according to any preceding claim further comprising from 0.05 to 15% by weight of a betaine.